

**Beeson Cosmetic Surgery**  
3660 Guion Road, Suite 320  
Indianapolis, IN 46222  
(317) 846-0846

Have these prescriptions filled several days before your surgery.

\_\_\_ Zovirax Capsules: This tablet will help prevent fever blisters and should be taken five capsules four times per day starting the day before surgery until the prescription is empty.

\_\_\_ Keflex: This is an antibiotic and should be taken one pill four times a day beginning the day before surgery and every day afterwards until the prescription is empty. Do not take day of surgery.

\_\_\_ Valium: Take one tablet an hour before bedtime the night before surgery and two tables the morning of surgery before \_\_\_\_\_.

\_\_\_ Novafed: This pill is to help decrease nasal congestion. Take one pill every twelve hours beginning the day after surgery.

\_\_\_ Percocet: Take one or two tablets every four to six hours as needed for pain after surgery.

\_\_\_ Compazine: This is a suppository and is for relief of nausea and vomiting. Use one suppository every six hours as needed after surgery.

\_\_\_ Theragran-M Vitamins/Vitamin C (1000mg): These vitamins will help to aid in the healing process. Take one of each vitamin each day beginning two weeks before surgery and every day afterwards for one month.

\_\_\_ Other: \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

\_\_\_ Hydrogen Peroxide  
\_\_\_ Q-Tips  
\_\_\_ ½ Inch Paper Tape  
\_\_\_ Baby Shampoo  
\_\_\_ Vaseline

\_\_\_ Bacitracin  
\_\_\_ 2 x 2 Gauze Squares  
\_\_\_ Crisco  
\_\_\_ Headband